

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS:

For control of Norway rats, roof rats and house mice.

URBAN AREAS: This product may be used in and around the periphery of homes, industrial, commercial and public buildings. This product may also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. This product may also be used in alleys. Do not use in sewers.

NON-URBAN AREAS: This product may be used inside of homes and agricultural buildings.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

HAWK[®]

RODENT BLOCK

KILLS RATS AND MICE

Kills Warfarin Resistant Norway Rats

Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

ACTIVE INGREDIENT: Bromadiolone;	
3-[3-(4'-Bromo-[1,1'-biphenyl]-4-yl)-3-hydroxy-1-	
phenylpropyl]-4-hydroxy-2H-1-benzopyran-2-one.....	
	0.005%
INERT INGREDIENTS*.....	
	99.995%
TOTAL	
	100.000%

*Contains Denatonium Benzoate

KEEP OUT OF REACH OF CHILDREN CAUTION

(See left panel for additional precautions.)

May be harmful or fatal if swallowed or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Wash hands after handling bait. If swallowed, call a physician at once.

For 24-hour emergency medical advice, call
1-877-854-2494

NET WEIGHT 10 lbs. (4.5 Kg)

Manufactured for:

Motomco Ltd.
Clearwater, FL 33755 U.S.A.

EPA EST. NO. 12455-WI-1

EPA REG. NO.12455-34-3240

DIRECTIONS FOR USE (Continued from right side panel)

APPLICATION DIRECTIONS:

Each HAWK RODENT BLOCK consists of a one-pound slab scored to be broken into four 4 oz. pieces.

RATS: Break and apply 1 to 4 pieces of HAWK RODENT BLOCK (usually at intervals of 15 to 30 feet) per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

MICE: Obtain 1 oz. bait pieces by breaking 4 oz. pieces of HAWK RODENT BLOCK in half and breaking each half in half. Use one 1 oz. piece of HAWK RODENT BLOCK per bait placement. Space placements at 8 to 12 foot intervals. Two 1 oz. pieces may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead exposed animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store only in original container, in a dry area inaccessible to children and pets.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Dispose of empty carton in a sanitary landfill or by incineration. If burned, stay out of smoke.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. If swallowed, call a physician at once.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K₁ intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds.