

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Keep out of reach of children and pets. Wear gloves when applying bait. With detergent and hot water, wash all implements used for applying bait. Do not use these implements for mixing, holding or transferring food or feeds. Treated baits should be placed in locations not accessible to nontarget wildlife and domestic animals, or in tamper-resistant bait boxes.

If swallowed by human beings, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of Vitamin K<sub>1</sub> combined with blood transfusions is indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin (Dicumarol).

**ENVIRONMENTAL HAZARDS**

This product is toxic to wildlife and fish. Use with care when applying in areas frequented by wildlife or adjacent to any body of water. Treated bait exposed on soil surface is hazardous to birds and other wildlife. Keep out of lakes, ponds, and streams.

**DIRECTIONS FOR USE**

**GENERAL CLASSIFICATION**

**FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA**  
It is a violation of State and Federal laws to use this product in a manner inconsistent with its labeling. (See Baiting Procedures.)

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.  
**STORAGE:** Store only in original container in a dry area inaccessible to children and pets.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty carton into application equipment. Then dispose of empty carton in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**IMPORTANT**

A single feeding of this anticoagulant bait will not control rodents. Bait must be eaten at several feedings on five or more successive days with no periods longer than 48 hours between feedings.

Keep this product in a closed container, free from odors which may contaminate the bait and reduce acceptability.

# TOMCAT

## GROUND SQUIRREL BAIT

ACTIVE INGREDIENT: Diphacinone; (2-Diphenylacetyl-1,3-indandione) .....	0.01%
INERT INGREDIENTS:.....	99.99%
TOTAL	100.0%

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

**FIRST AID TREATMENT**

**IF SWALLOWED:** Call a physician or Poison Control Center; drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

For 24-hour emergency medical advice, call  
1-877-854-2494

NET CONTENTS: 50 lbs.

EPA EST. NO. 12455-WI-1      EPA SLN NO. CA 780146  
CALIF. REG. NO. 12455-50003ZA

Manufactured by:



**Bell Laboratories, Inc.**

Madison, WI 53704

Distributed By:

**Motomco Ltd.**

Madison, WI 53704 U.S.A.

### BAITING PROCEDURES

**NOTE:** Do not use in a manner that will contaminate feed or foodstuff.

**GROUND SQUIRRELS, *Spermophilus beecheyi*, *S. belding***-Scatter by hand approximately 2 oz. of bait (one handful) evenly over 40 to 50 square feet near active burrows or runways. Retreat every other day for 3 to 4 applications.

An uninterrupted supply of bait should be available for 6 to 8 days. Don't pile bait. The scattering of bait takes advantage of the squirrels' natural foraging habits and limits domestic livestock and wildlife from picking it up.

**DEER MICE, *Peromyscus spp.***- Deer Mouse Control in Forests: Spread bait evenly by hand, mechanical spreader or aircraft at the rate of two pounds per acre through the area where seeding will later take place. Sufficient bait should be applied to last the mouse population a minimum of four days (preferably without rain). Bait application should precede the sowing of conifer seed by 10 days to 2 weeks.

**HOUSE MICE, *Mus musculus***- Place tablespoon amounts (1/4 to 1/2 ounce) of bait in bait box or shallow container, preferably in protected feeder station. Place bait station at 8- to 12- foot intervals in dry locations such as in concealed places, in corners, or along walls where house mice feed, drink or frequent. Inspect stations daily and add bait as needed; increase the amount when bait in feeder is entirely consumed overnight. Replace moldy or old bait with fresh bait. The baits being replaced should immediately be disposed of in such manner to prevent ingestion by children, pets, wildlife and domestic animals. An uninterrupted supply of bait should be maintained as long as any bait is taken which may be 2 to 4 weeks.

Where continuous source of infestation is present, permanent bait stations should be established and the bait replenished as needed. Bait should be picked up and disposed of upon termination of control program.

**MEADOW MICE, *Microtus californicus*, *M. montanus*** - Lightly scatter tablespoon amounts (1/4 to 1/2 ounce) of bait near active burrows or in runways. Repeat treatment every other day for three treatments.

**WARRANTY:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.