



WITH

CONTAINS LUMITRACK®

For Use In and Around Agricultural Buildings Only

KILLS RATS, MICE & MEADOW VOLES*

Net Weight: 16 lbs (7.2kg)

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panels for First Aid and additional precautionary statements.

*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat

FIRST AID

HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE

If Swallowed

- Call a poison control center, doctor, or 1-877-854-2494, or 1-800-858-7378** immediately for treatment advice.
- Have a person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- **Also call this number for information on health concerns and pesticide incidents.

NOTE TO PHYSICIAN

If swallowed, this material may reduce the dotting ability of the blood and cause bleeding. It ingested, administer Vitamin K₁ intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times

TREATMENT FOR PET POISONING

If animal eats bait, call veterinarian at once.

NOTE TO VETERINARIAN

Anticoagulant Brodifacoum: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin k_1 at doses of 2–5 mg/kg. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets, or nontarget animals to rodenticides. To help to prevent exposure:

- 1. Store unused product out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container and unused, spoiled, or unconsumed bait as specified on this label.

Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used wherever children, pets, non-target mammals, or birds may have access to the bait placement location.

USE RESTRICTIONS: This product may only be used to control the following rodent pests: House mouse (Mus musculus), Norway rat (Rattus norvegicus), Roof rat (Rattus rattus), Cotton rat* (Sigmodon hispidus), Eastern harvest mouse* (Reithrodontomys humuli), Golden mouse* (Ochrotomys nuttalli), Polynesian rat* (Rattus exulans), Meadow vole* (Microtus pennsylvanicus), White-throated woodrat* (Neotoma albigula), Southern plains woodrat* (Neotoma micropus), and Mexican woodrat* (Neotoma mexicana). This product must be used in and within 100 feet of agricultural buildings and man-made agricultural structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Fence and perimeter baiting, beyond 100 feet from an agricultural structure as defined above, is prohibited. This product must not be applied directly to food or feed crops.

Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant ball stations. Do not broadcast ball.

MOTOMCO FARM & RANCH™



CONTAINS **LUMITRACK**®

For Use In and Around Agricultural Buildings Only

KILLS RATS, MICE & MEADOW VOLES'

KILLS WARFARIN RESISTANT NORWAY RATS & HOUSE MICE

Norway rats, roof rats and house mice may consume a lethal dose in one night's feeding with first dead rodents appearing five or six days after feeding begins.

 ACTIVE INGREDIENT:
 0.0025%

 Brodifacoum (CAS #56073-10-0):
 0.0025%

 OTHER INGREDIENTS†:
 99.9975%

 †Contains Denatonium Benzoate
 TOTAL

 100.0000%

KEEP OUT OF REACH OF CHILDREN CAUTION

See side panels for First Aid and additional precautionary statements.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

Pesticide Disposal: Wastes resulting from the use of this product may be placed in trash or delivered to an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. [Plastic:] Offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or by incineration. In most states, burning is not allowed. [Paper:] Dispose of empty container by placing in trash, at an approved waste disposal facility or by incineration. In most states, burning is not allowed.

WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



3699 Kinsman Blvd. Madison, WI 53704 U.S.A. www.motomco.com

The World Leader In Rodent Control Technology®

Net Weight: **16 lbs** (7.2kg)

EPA REG. NO. 12455-139-3240 EPA EST. NOs. 12455-WI-1k, 12455-WI-2P, 12455-WI-3w Superscript is first letter of the lot number. **DIRECTIONS FOR USE** (Continued from other panel)

Do not sell this product in individual containers holding less than 8 pounds of bait.

Burrow baiting with JAGUAR® Soft Bait is prohibited.

SELECTION OF TREATMENT AREAS: Determine areas where rats, mice, or meadow voles* will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in corners and concealed places, between floors and walls, or in locations where rats, mice, or meadow voles* or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

RATS: Place 6 to 28 sachets at intervals of 15 to 30 feet per placement in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

MICE AND MEADOW VOLES*: Place 1 sachet per placement. Space placements at 8 to 12 foot intervals in infested areas. Three sachets may be needed at points of very high activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse or meadow vole* activity cease.

Bait may be removed from sachet and placed in a feeding device that secures the unwrapped bait in a bait station. Discard empty wrapper in the trash.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Wearing gloves, collect and dispose of all dead, exposed animals and leftover bait. To prevent reinfestation, limit sources of rodent food, water, and

harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Keep away from children, domestic animals and pets.

All handlers (including applicators) must wear shoes plus socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

User Safety Requirements


Follow manufacturer's instruction for deaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into dean dothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to mammals and birds. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff also may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate.

*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat

063017/11-15



•587JS-2

JAGUAR®SOFT BAIT

KEEP OUT OF REACH OF CHILDREN CAUTION DIRECTIONS FOR USE It is a violation of Federal law to use this product in a

manner inconsistent with its labeling. See outer package label for complete DIRECTIONS FOR USE including USE RESTRICTIONS.

NET WEIGHT: 0.53 oz (15 g)

EPA REG. NO. 12455-139-3240
EPA EST. NOs. 12455-WI-1^k, 12455-WI-2^k, 12455-WI-3^w
Superscript is first letter of the lot number.

INDIVIDUAL SALE IS PROHIBITED BY LAW

021418/11-15

2607-1