

# Мотрмсо



MADE IN USA

# ROUGH CUT NUGGETS

# KILLS RATS, MICE & MEADOW VOLES\*

# Net Weight: 4 lbs (1.8 kg)

Diphacinone (CAS #82-66-6): . . . . . . . 0.005% OTHER INGREDIENTS: . 99.995% TOTAL 100.000%

**KEEP OUT OF REACH OF CHILDREN** CAUTION

See back panels for First Aid and additional precautionary statements

EPA REG. NO. 12455-81-3240 EPA EST. NOs. 12455-WI-1<sup>k</sup>,12455-WI-2<sup>p</sup>, 12455-WI-3<sup>w</sup> Superscript is first letter of the lot number

### DIRECTIONS FOR USE

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this

#### IMPORTANT:

WITH

ACINO

Do not expose children, pets, or nontarget animals to rodenticides. To help to prevent accidents: 1. Store unused product out of reach of children and pets.

2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potential destructive animals, or in areas prior to vandalism. 3. Dispose of product container and unused, spoiled, or unconsumed bait as specified on this label.

Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used wherever children, pets, non-target mammals, or birds may have access to the bait placement location.

USE RESTRICTIONS: This product may only be used to control the following rodent pests in and around man-made structures: House mouse (Mus musculus), Norway rain and advergicus), Roof rat (Rattus rattus), Cotton rat' (Signodon hispidus), Eastern harvest mouse\* (Reithrodontomys humuli), Golden mouse\* (Ochrotomys nuttalli), Polynesian rat (Rattus exulans), Meadow vole' (Microtus pennsylvanicus) White-throated woodrat\* (Neotoma albigula), Southem plains woodrat\* (Neotoma micropus), and Mexican woodrat<sup>\*</sup> (*Neotoma mexicana*). This product must be used and within 100 feet of man made structures constructed in a manner co as to be **KEEP OUT OF REACH OF CHILDREN** in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains acrarth, docks and port or terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops.

his product may be applied to active rodent burrows to control rats within or beyond 100 feet of buildings and man-made structures, provided that infestations of these rodents have been confirmed. Bait must be placed no less than 6 inches into active rat burrows. Do not broadcast bait. Because rat infestations may occur in areas farther than 100 feet from buildings and man-made structures when the rodents have ample supplies of food and cover, efforts should be made to remove food trash, garbage, clutter and debris.

Do not place near or inside ventilation duct openings. Do not contaminate water, food, Do not place near or inside verification duct openings. Do not charameter when handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected indicated on the label. Buyer assumes all risk of use and/or handling of this product other than the label. WARRANTY: To the extent consistent with applicable law, seller makes not material when such use and/or handling is contrary to label instructions.

Do not sell this product in individual containers holding less than 4 pounds of bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats, mice or meadow voles\* will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as

#### APPLICATION DIRECTIONS:

infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

714

Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat



TO OPEN: TEAR TOWARD CENTER FROM BOTH NOTCHES

PRESS ZIPPER TO RESEAL

It is a violation of Federal law to use this product in a manner inconsistent with its

RATS: Apply 4 to 16 oz. of bait (usually at intervals of 15 to 30 feet) per placement in

041019/09-15



Мотрмсо

### ROUGH CUT NUGGETS

### KILLS RATS, MICE & MEADOW VOLES FOR USE IN AND AROUND HOMES AND BUILDINGS PELLETED BAIT

ACTIVE INGREDIENT: Diphacinone (CAS #82-66-6): . . . . . . 0.005% OTHER INGREDIENTS: . . . . . . . . . . . . . . 99.995% τοται 100.000%

# CAUTION

See side panels for First Aid and additional precautionary statements

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposa esticide Storage: Store only in original container in a cool, dry place naccessible to children and pets. Keep containers closed and away from other chemicals.

Pesticide Disposal: Dispose of wastes resulting from the use of this product in trash or at an approved waste disposal facility. Container Handling: Nonrefillable container. Do not reuse or refill this container. [Plastic:] Offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or by incineration. In most states, burning is not allowed.

> EPA REG. NO. 12455-81-3240 EPA EST. NOs. 12455-WI-1k, 12455-WI-2p, 12455-WI-3w Superscript is the first letter of the lot number.

### Net Weight: **4 lbs** (1.8 kg)

Available Exclusively From:

Мотрмсо www.m MADE IN USA





#### APPLICATION DIRECTIONS (Continued from other panel)

MICE AND MEADOW VOLES\*: Apply 1/4 to 1/2 oz. (1 to 2 level tablespoons) of bait at 8 to 12 foot intervals in infested areas. Larger placements (up to 2 oz.) may be needed at points of very high activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse or meadow vole\* activity cease.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Using waterproof gloves, collect and dispose of all dead, exposed animals and leftover bait. To prevent reinfestation limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

#### PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals CAUTION: Harmful if swallowed. Keep away from children, domestic animals and

pets. All handlers (including applicators) must wear: shoes plus socks, and gloves. Any person who retrieves carcasses or unused bait following application of this product nust wear waterproof gloves.

User Safety Requirements Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash induct. Wash inducted of gloves before tentoring. As soon as possible, wash ands thoroughly after applying bait and before eating, drinking, chewing gum using tobacco or using the toilet and change into clean clothing.

FIRST AID HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE

- Call a poison control center, doctor, or 1-877-854-2494
- immediately for treatment advice.
- Have a person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by the poison contro center or doctor.

#### NOTE TO PHYSICIAN

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.




Anticoagulant Diphacinone: For animals ingesting bait and/or showin poisoning signs (bleeding or elevated prothrombin times), give Vitamin K1.

#### ENVIRONMENTAL HAZARDS

This product is extremely toxic to mammals and birds and other wildlife. Dogs, cats and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

\*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat





3699 Kinsman Blvd. Madison, WI 53704 U.S.A.

The World Leader In Rodent Control Technology®